



FORM PTO 1390 (REV 5-93)		US DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY DOCKET NUMBER 2001_1838A	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. §371				U.S. APPLICATION NO. (if known, insert OFFICE) NEW 10/009962	
International Application No. PCT/JPO0/03806		International Filing Date June 12, 2000		Priority Date Claimed June 14, 1999	
Title of Invention PLANT THERMOGENIC GENES AND PROTEINS					
Applicant(s) For DO/EO/US Kikukatsu ITO					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. §371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. §371. 3. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1). 4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. §371(c)(2)) a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US) 6. <input checked="" type="checkbox"/> A translation of the International Application into English (35 U.S.C. §371(c)(2)). ATTACHMENT A 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3)). a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19. 9. <input checked="" type="checkbox"/> An unexecuted oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)). ATTACHMENT B 10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)). Items 11. to 14. below concern other document(s) or information included: 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. ATTACHMENT C 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> Other items or information:					


THE COMMISSIONER IS AUTHORIZED
TO CHARGE ANY DEFICIENCY IN THE
FEE FOR THIS PAPER TO DEPOSIT
ACCOUNT NO. 23-0975.

U.S. APPLICATION NO. (if known, see 37 CFR 1.5) NEW		INTERNATIONAL APPLICATION NO. PCT/JP00/03806		ATTORNEY'S DOCKET NO. 2001 1838A					
15. [X] The following fees are submitted BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee nor international search fee paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 International Search Report has been prepared by the EPO or JPO \$ 890.00 International preliminary examination fee not paid to USPTO but international search paid to USPTO \$ 740.00 International preliminary examination fee paid to USPTO but claims did not satisfy provisions of PCT Article 33(1)-(4) \$ 690.00 International preliminary examination fee paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$ 100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <th style="width:70%;">CALCULATIONS</th> <th style="width:30%;">PTO USE ONLY</th> </tr> <tr> <td style="height: 100px; vertical-align: bottom;">\$890.00</td> <td></td> </tr> </table>		CALCULATIONS	PTO USE ONLY	\$890.00	
CALCULATIONS	PTO USE ONLY								
\$890.00									
Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$					
Claims	Number Filed	Number Extra	Rate						
Total Claims	-20 =		X \$18.00	\$					
Independent Claims	4 - 3 =	1	X \$84.00	\$84.00					
Multiple dependent claim(s) (if applicable)				+ \$280.00					
TOTAL OF ABOVE CALCULATIONS =				\$974.00					
[] Small Entity Status is hereby asserted. Above fees are reduced by 1/2.				\$					
SUBTOTAL =				\$974.00					
Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+ \$					
TOTAL NATIONAL FEE =				\$974.00					
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40 per property				+ \$					
TOTAL FEES ENCLOSED =				\$974.00					
				Amount to be refunded \$					
				Amount to be charged \$					
a. [X] A check in the amount of \$974.00 to cover the above fees is enclosed. A duplicate copy of this form is enclosed. b. [] Please charge my Deposit Account No. 23-0975 in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. [] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-0975.									
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.									
19. CORRESPONDENCE ADDRESS <div style="text-align: center;">  000513 PATENT TRADEMARK OFFICE </div>			By:  Warren M. Cheek, Jr., Registration No. 33,367 WENDEROTH, LIND & PONACK, L.L.P. 2033 "K" Street, N.W., Suite 800 Washington, D.C. 20006-1021 Phone: (202) 721-8200 Fax: (202) 721-8250 December 14, 2001						

JG09 Rec'd PCT/PTO 1 4 DEC 2001

FORM PTO 1390 (REV 5-99)		US DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY DOCKET NUMBER 2001_1838A
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. §371				U.S. APPLICATION NO. (if known, see 37 CFR 1.53) NEW 10/009962
International Application No. PCT/JP00/03806		International Filing Date June 12, 2000		Priority Date Claimed June 14, 1999
Title of Invention PLANT THERMOGENIC GENES AND PROTEINS				
Applicant(s) For DO/EO/US Kikukatsu ITO				
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:				
1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. §371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. §371. 3. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. §371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. §371(b) and PCT Articles 22 and 39(1). 4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. §371(c)(2)) a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau. c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US) 6. <input checked="" type="checkbox"/> A translation of the International Application into English (35 U.S.C. §371(c)(2)). ATTACHMENT A 7. <input type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. §371(c)(3)). a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19. 9. <input checked="" type="checkbox"/> An unexecuted oath or declaration of the inventor(s) (35 U.S.C. §371(c)(4)). ATTACHMENT B 10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. §371(c)(5)). Items 11. to 14. below concern other document(s) or information included: 11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. ATTACHMENT C 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> Other items or information:				

THE COMMISSIONER IS AUTHORIZED
 TO CHARGE ANY DEFICIENCY IN THE
 FEE FOR THIS PAPER TO DEPOSIT
 ACCOUNT NO. 23-0975.

U.S. APPLICATION NO. 107009962 NEW		INTERNATIONAL APPLICATION NO. PCT/JP00/03806		ATTORNEY'S DOCKET NO. 2001 1838A					
15. <input checked="" type="checkbox"/> The following fees are submitted BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Neither international preliminary examination fee nor international search fee paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00 International Search Report has been prepared by the EPO or JPO \$ 890.00 International preliminary examination fee not paid to USPTO but international search paid to USPTO \$ 740.00 International preliminary examination fee paid to USPTO but claims did not satisfy provisions of PCT Article 33(1)-(4) \$ 690.00 International preliminary examination fee paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$ 100.00 ENTER APPROPRIATE BASIC FEE AMOUNT =				<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <th style="width:50%;">CALCULATIONS</th> <th style="width:50%;">PTO USE ONLY</th> </tr> <tr> <td style="height: 100px; vertical-align: bottom;">\$890.00</td> <td></td> </tr> </table>		CALCULATIONS	PTO USE ONLY	\$890.00	
CALCULATIONS	PTO USE ONLY								
\$890.00									
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width:50%;">\$</td> <td style="width:50%;"></td> </tr> </table>		\$			
\$									
Claims	Number Filed	Number Extra	Rate						
Total Claims	-20 =		X \$18.00	\$					
Independent Claims	4 - 3 =	1	X \$84.00	\$84.00					
Multiple dependent claim(s) (if applicable)			+ \$280.00	\$					
TOTAL OF ABOVE CALCULATIONS =				\$974.00					
<input type="checkbox"/> Small Entity Status is hereby asserted. Above fees are reduced by 1/2.				\$					
SUBTOTAL =				\$974.00					
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	\$				
TOTAL NATIONAL FEE =				\$974.00					
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40 per property +				\$					
TOTAL FEES ENCLOSED =				\$974.00					
				Amount to be refunded	\$				
				Amount to be charged	\$				
a. <input checked="" type="checkbox"/> A check in the amount of \$974.00 to cover the above fees is enclosed. A duplicate copy of this form is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. 23-0975 in the amount of \$_____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-0975.									
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.									
19. CORRESPONDENCE ADDRESS <div style="text-align: center;">  000513 PATENT TRADEMARK OFFICE </div>			By: <u>Warren M. Cheek, Jr.</u> Warren M. Cheek, Jr., Registration No. 33,367 WENDEROTH, LIND & PONACK, L.L.P. 2033 "K" Street, N.W., Suite 800 Washington, D.C. 20006-1021 Phone: (202) 721-8200 Fax: (202) 721-8250 December 14, 2001						



Rec'd PCT/PTO 13 MAY 2002

#7/a

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of : Confirmation No. 3213
Kikukatsu ITO : Docket No. 2001-1838A
Serial No. 10/009,962 : Group Art Unit Not Yet Assigned
Filed January 23, 2002 : Examiner Not Yet Assigned
PLANT THERMOGENIC GENES :
AND PROTEINS :

IF CLAIMS ARE NOT AUTHORIZED
TO CHARGE ANY DEFICIENCY IN THE
FEES FOR THIS PAPER TO DEPOSIT
ACCOUNT NO. 23-0975

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents,
Washington, D.C. 20231

Sir:

Responsive to the Notice dated February 13, 2002, the time for filing thereto being extended for one month in accordance with the Petition for Extension submitted concurrently herewith, please amend the above-identified application as follows:

In the Specification:

Page 1, line 1, delete the entire heading.

between lines 3 and 6, insert the following new heading:

Background of the Invention

line 6, replace the heading with the following new heading:

1. Field of the Invention

line 16, replace the heading with the following new heading:

2. Description of the Related Art

[illegible]

However, the expression of StUCP and AtPUMP have been induced by low temperature.

Therefore, it has been suggested that these genes are involved in heat production (Laloi et al., 1997; Maia et al., 1998).

line 26, replace the heading with the following new heading:

Summary of the Invention

Page 4, replace the paragraph beginning at line 23 with the following paragraph:

Fig. 3 compares the alignment of amino acid sequences of SfUCPA (SEQ ID No. 2) and SfUCPB (SEQ ID No. 4), together with potato UCP (StUCP) (SEQ ID No. 5), *Arabidopsis* UCP (AtPUMP) (SEQ ID No. 6) and human UCP (human UCP 1, 2 and 3 corresponding to SEQ ID Nos. 7, 8 and 9, respectively). The asterisks (*) attached under the sequences indicate the same amino acid sequence, and the dot (.) indicates the conservative change in all of the sequences. The boldface indicates the same sequence between SfUCPA (SEQ ID No. 2) and SfUCPB (SEQ ID No. 4). The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. The characteristic domains of energy transfer proteins typical of mitochondria are surrounded by a square. The shaded bars (I~VI) above the upper sequence show estimated transmembrane domains.

Page 5, line 25, replace the heading with the following new heading:

Description of the Preferred Embodiments

Page 10, replace the paragraph beginning at line 28 with the following paragraph:

The total RNA was extracted from the spadix of skunk cabbage (*Symplocarpus foetidus*) and the complete RNA was determined on 1.0% agarose gel electrophoresis (Ito et al., 1999). Using a mRNA isolation kit (Pharmacia), a clone associated with the UCP gene family was isolated from the purified poly(A)⁺RNA by RT-PCR. The first strand cDNA was prepared by annealing 20 pmol of cDNA primed primer (5'-TTTTTTTTTTTTTTTTTTTTTTTTTTT-3') (SEQ ID No. 10) into poly(A)⁺RNA (0.1 µg), followed by extension with 10 units of reverse transcriptase (New England Biolab) at 37°C for 30 minutes in 20 µl of 1xRT buffer containing 10mM 1,4-dithiothreitol and 0.5mM dNTP. The composition of the reaction solution is as follows.

- 10mM Tris-HCl (pH 8.0);
- 50mM KCl;
- 1.5mM MgCl₂;
- 4mM dNTP;
- 0.2 unit of EX Taq polymerase (Takara); and
- 10pmol of two degenerate primers corresponding to the conserved amino acid

sequence of the UCP family:

ZF1 (5'-CCIYTIGAYACIGCIAAR-3') (SEQ ID No. 11)

ZR1 (5'-ACWTTCCAISYICCIAWIC-3') (SEQ ID No. 12).

Page 13, line 15, delete the entire heading.

[illegible]

What is claimed is:

Page 17, line 1, replace the heading with the following new heading

In the Sequence Listing:

Please replace the Sequence Listing of record with the attached substitute Sequence

The foregoing amendments are presented to place the application in compliance with the sequence rules under 37 CFR 1.821-1.825.

Applicants have submitted a Sequence Listing in both paper and computer readable form

Additional amendments to the specification have also been effected to put the specification

conformance with U.S. practice. Also, the additional sequences disclosed in the specification and Figure 3 have been identified and labeled as required under U.S. practice.

With regard to the Notice also requesting that an executed Oath and Declaration of the Inventors needs to be submitted, Applicants wish to note that an executed Oath and Declaration was submitted on January 23, 2002. A copy of the submitted executed Declaration is enclosed herewith along with the cover letter (indicating the filing of the executed Declaration). Applicants respectfully request that the Patent Office review the application papers to ensure that the executed Declaration is present in the file.

Attached hereto is a marked-up version of the changes made to the specification by the current amendment. The attached page is captioned "Version with markings to show changes made."

In view of the foregoing, it is believed that each requirement set forth in the Notice has been satisfied, and that the application is now in compliance with the sequence rules under 37 CFR 1.821-1.825. Accordingly, favorable examination on the merits is respectfully requested.

Respectfully submitted,

Kikukatsu ITO

By: 

Lee Cheng
Registration No. 40,949
Attorney for Applicant

LC/gtn
Washington, D.C. 20006-1021
Telephone (202) 721-8200
Facsimile (202) 721-8250
May 13, 2002



DESCRIPTION

Plant Thermogenic Genes and Proteins

5

Background of the Invention
1. Technical Field of the Invention

10

The present invention relates to plant thermogenic genes and proteins. More particularly, the invention relates to thermogenic genes derived from a skunk cabbage (*Symplocarpus foetidus*) and gene products (proteins). Those genes and proteins are useful in breeding of cold-avoidance plants, medical treatment of diabetes mellitus or obesity, or development of novel thermogenic bio-materials.

15

2. Description of the Related Art
Background-

20

25

Stresses due to low temperatures, droughts and salinity are common harmful environmental factors that terrestrial plants encounter. Among these stresses, it has been considered that cellular injury due to the low temperature is the most important factor which restricts productivity of crops (Levitt, 1980). To resist the low temperature stress, cold-hardy plants such as wheat or rye have a variety of physiological and metabolic responses which lead to cold acclimation (Sakai and Larcher, 1987; Steponkus, 1984; Thomashow, 1998; Uemura and Steponkus, 1997). In contrast, it is known that some plants including skunk cabbage have a specialized system by which the plants generate heat to avoid freezing (Knutson, 1974; Nagy et al., 1972; Schneider and Buchenen, 1980).

30

The temperature of the flower in the spadix of skunk cabbage, which flowers in early spring, has been known to maintain its temperature at higher

the C-terminal region and increases by free fatty acids (Jezek et al., 1998; Lin and Klingenberg, 1982; Katiyar and Shrago, 1989; Rial et al., 1983; Sluse et al., 1998).

On the contrary, 2 cDNAs encoding UCP-like proteins of plant origin were
 5 isolated from potato (StUCP: Laloi et al., 1997) and from *Arabidopsis* (AtPUMP: Maia et al., 1998). Since the expression of StUCP was mainly detected in the flower and the fruit, it has been postulated that StUCP may concern respiration during flowering and maturation of the fruit together with the AOX activity (Laloi et al., 1997).

10

Potato and *Arabidopsis* have been considered to be non-thermogenic plants. However, the expression of StUCP and AtPUMP ^{have been} induced by low temperature. Therefore, it has been suggested that these genes are involved in ~~the~~ heat production (Laloi et al., 1997; Maia et al., 1998).

15

In the thermogenic plants such as skunk cabbage, however, UCP-mediated thermoigenic mechanisms have not yet been identified.

The purpose of the invention of this application is to provide unidentified
 20 novel UCP genes derived from a thermogenic plant, skunk cabbage.

The additional purpose of this application is to provide skunk cabbage UCPs which are expression products of the novel genes.

25

Summary of the **Disclosure of Invention**

The invention provides thermogenic genes derived from skunk cabbage, i.e., gene SfUCPa of which cDNA comprises the base sequence of SEQ ID NO: 1,
 30 and gene SfUCPb of which cDNA comprises the base sequence of SEQ ID NO: 3.

Moreover, the invention provides thermogenic proteins, i.e., protein SfUCPA expressed from SfUCPa, which comprises the amino acid sequence of SEQ ID NO: 2, and protein SfUCPB expressed from SfUCPb, which comprises the amino acid sequence of SEQ ID NO: 4.

10

In addition, the invention provides cDNA having the base sequence of SEQ ID NO: 1 or a partial sequence thereof, and cDNA having the base sequence of SEQ ID NO: 3 or a partial sequence thereof.

Brief Description of Drawings

Fig. 1 shows the change of the temperature of the spadix in skunk cabbage and that of ambient temperature with a lapse of time.

Fig. 2 shows the results of northern blotting, indicating the expression profile of SfUCPa (A) and SfUCPb (B) in the spadix and leaf of skunk cabbage at room temperature (RT) and during cold treatment (4°C for 3 days). The lower figures respectively show the results of ethidium bromide staining of non-decomposed rRNA.

Fig. 3 compares the alignment of amino acid sequences of SfUCPA^(SEQ ID No. 2) and SfUCPB^(SEQ ID No. 4), together with potato UCP (StUCP)^(SEQ ID No. 5), *Arabidopsis* UCP (AtPUMP)^(SEQ ID No. 6) and human UCP^(SEQ ID No. 7). The asterisks (*) attached under the sequences indicate the same amino acid sequence, and the dot (.) indicates the conservative change in all of the sequences. The boldface indicates the same sequence between SfUCPA^(SEQ ID No. 2) and SfUCPB^(SEQ ID No. 4). The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. The characteristic domains of energy transfer proteins typical of mitochondria are

surrounded by a square. The shaded bars (I-VI) above the upper sequence show estimated transmembrane domains.

Fig. 4 shows a hydrophobic plot of SfUCPA. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM6.

Fig. 5 shows a diagrammatic illustration of SfUCPA topology in the mitochondria membrane.

10

Fig. 6 shows a hydrophobic plot of SfUCPB. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM4 and TM6.

15

Fig. 7 shows a diagrammatic illustration of SfUCPB topology in the mitochondria membrane.

Fig. 8 shows the results of *in vitro* translation using respective cDNAs of the genes SfUCPa and SfUCPb as templates. (-) indicates a control, S a sense RNA, and AS an antisense RNA. The asterisk (*) indicates a non-specific product and the empty circle denotes the position of a low molecular translated artificial product synthesized from a small ORF.

25

Description of the Preferred Embodiments **Best Mode for Carrying Out the Invention**

In the gene SfUCPa of the present invention, its cDNA has the base sequence of SEQ ID NO: 1 and encodes the protein SfUCPA having the amino acid sequence of SEQ ID NO: 2, of which the estimated molecular weight is 32.6 kDa.

30 In the gene SfUCPb of the present invention, its cDNA (SEQ ID NO: 3) encodes

many include mammal cultured cells such as monkey renal cell COS7, Chinese hamster ovarian cell CHO, etc., budding yeast, fission yeast, silkworm cell, *Xenopus* egg cell, and the like are commonly used, but not limited thereto. In order to introduce the expression vector into eucaryotic cells, a known method
5 such as electroporation, calcium phosphate method, liposome method, DEAE dextran method, and the like can be utilized.

After expression of the proteins in procaryotic cells or eucaryotic cells according to the aforementioned method, the desired proteins are isolated and
10 purified from the culture in the known combined procedures for separation. For example, treatment with a denaturant (e.g., urea) or surface activator, ultrasonication, digestion with enzymes, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity
15 chromatography, reverse phase chromatography, and the like are involved.

The proteins of the invention, SfUCPA and SfUCPB, also include peptide fragments (5 amino acids or more) involving the optional partial amino acid sequences of SEQ ID NOS: 2 and 4. In addition, the proteins of the invention also
20 include fusion proteins with other optional proteins.

The following examples serve to illustrate the invention of this application specifically in more detail, but are not intended to limit the scope of the invention.

25

Example 1: Cloning of cDNA

The total RNA was extracted from the spadix of skunk cabbage (*Symplocarpus foetidus*) and the complete RNA was determined on 1.0% agarose
30 gel electrophoresis (Ito et al., 1999). Using a mRNA isolation kit (Pharmacia), a

clone associated with the UCP gene family was isolated from the purified poly(A)⁺RNA by RT-PCR. The first strand cDNA was prepared by annealing 20 pmol of cDNA primed primer (5'-TTTTTTT/TTTTTTT/TTTTTTT/TTTTTTT-3') into poly(A)⁺RNA (0.1μg), followed by extension with 10 units of reverse transcriptase (SEQ ID NO. 10) (New England Biolab) at 37°C for 30 minutes in 20μl of 1×RT buffer containing 10mM 1,4-dithiothreitol and 0.5mM dNTP. The composition of the reaction solution is as follows.

- 10mM Tris-HCl (pH 8.0);
- 50mM KCl;
- 1.5mM MgCl₂;
- 4mM dNTP;
- 0.2 unit of EX Taq polymerase (Takara); and
- 10pmol of two degenerate primers corresponding to the conserved amino acid sequence of the UCP family:

ZF1 (5'-CCCTTIGAYACIGCIAAR-3') (SEQ ID NO. 11)

ZR1 (5'-ACWTTGCCAISYICCLAWIC-3') (SEQ ID NO. 12)

PCR cycle was carried out as follows.

(94°C: 0.5 minute; 50°C: 1 minute; 72°C: 1 minute)×35

Among the PCR products obtained in the above method, the amino acid sequence estimated from the sequence of about 0.8kb cDNA fragment had very high homology to one of the reading frame sequences of the UCP gene family. This fragment, accordingly, was cloned into T-vector (clone p2-1) and used as a probe for library screening.

cDNA (5μg) prepared from the spadix was inserted into λgt11 phage according to the known method (Sambrook et al., 1989) to construct a cDNA library. From this library, 8 clones positive to the above-described probes were isolated and sub-cloned into the pBluescript SK plasmid (Stratagene). From

linearized, on which a sense- or anti-sense RNA was transcribed with T7 RNA polymerase or T3 RNA polymerase according to the protocol of MAXICRIPT transcription kit (Ambion). An equal amount of RNA (4µg) was provided for in vitro translation reaction using a wheat germ extract (Promega) in the presence of ³⁵S-methionine (Amersham). The translation product was analyzed by SDS-PAGE. The gel was fixed, incubated in Amplify (Amersham), then dried, and fluorometrically analyzed.

As a result, it was confirmed that, as shown in Fig. 8, the initiation codon and the stop codon of cDNA isolated in Example 1 functioned successfully since a protein having an expected molecular weight was produced from any of cDNAs only when the sense RNA was used as a template.

~~Industrial Applicability~~

As described previously, this application provides novel thermogenic genes SfUCPa and SfUCPb as well as their gene products, i.e., thermogenic proteins SfUCPA and SfUCPB, derived from skunk cabbage (*Symplocarpus foetidus*), and cDNAs used for gene engineering mass production of these proteins. These genes and proteins allow development of low temperature-tolerant plants, development of drugs or methods for treatment of diabetes mellitus or obesity, or development of novel heat generating materials from plants.

References

- Berthold and Siedow (1993) Plant Physiol. 101, 113-119.
 Boss et al. (1997) FEBS Lett. 408, 39-42.
 Fleury et al. (1997) Nature Genetics 15, 269-272.

CLAIMS

What is claimed is:

1. A thermogenic gene SfUCPa derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 1.

5

2. A thermogenic gene SfUCPb derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 3.

3. A thermogenic protein SfUCPA expressed from the gene SfUCPa of Claim 1, which comprises the amino acid sequence of SEQ ID NO: 2.

10

4. A Thermogenic protein SfUCPB expressed from the gene SfUCPb of Claim 2, which comprises the amino acid sequence of SEQ ID NO: 4.

5. A DNA fragment comprising the base sequence of SEQ ID NO: 1 or a partial sequence thereof.

15

6. A DNA fragment comprising the base sequence of SEQ ID NO 3: or a partial sequence thereof.

20

ABSTRACT OF THE DISCLOSURE

The inventions of this application include thermogenic genes named SfUCPa and SfUCPb which are derived from skunk cabbage. cDNA of each gene
5 comprises the base sequence of SEQ ID NO: 1 and 3, respectively. Thermogenic proteins, SfUCPA and SfUCPB, are expressed from genes SfUCPa and SfUCPb, comprises the amino acid sequence of SEQ ID NO: 2 and 4.

8/PRTS

107009962
JC05 Rec'd PGT/PTO 14 DEC 2007

1

DESCRIPTION

Plant Thermogenic Genes and Proteins

5

Technical Field

The present invention relates to plant thermogenic genes and proteins. More particularly, the invention relates to thermogenic genes derived from a
10 skunk cabbage (*Symplocarpus foetidus*) and gene products (proteins). Those genes and proteins are useful in breeding of cold-avoidance plants, medical treatment of diabetes mellitus or obesity, or development of novel thermogenic bio-materials.

15

Background

Stresses due to low temperatures, droughts and salinity are common harmful environmental factors that terrestrial plants encounter. Among these
20 stresses, it has been considered that cellular injury due to the low temperature is the most important factor which restricts productivity of crops (Levitt, 1980). To resist the low temperature stress, cold-hardy plants such as wheat or rye have a variety of physiological and metabolic responses which lead to cold acclimation (Sakai and Larcher, 1987; Steponkus, 1984; Thomashow, 1998; Uemura and
25 Steponkus, 1997). In contrast, it is known that some plants including skunk cabbage have a specialized system by which the plants generate heat to avoid freezing (Knutson, 1974; Nagy et al., 1972; Schneider and Buchenen, 1980).

The temperature of the flower in the spadix of skunk cabbage, which
30 flowers in early spring, has been known to maintain its temperature at higher

than +10°C even when the ambient temperature falls to -15°C (Knutson, 1974). For example, thermoscopic analysis using infrared camera indicates homeothermic behavior of the surface temperature of the spadix (Fig. 1). It should be noted that, in this experiment, the plants were placed in the growth chamber
 5 and the air temperature was gradually decreased. As clearly seen from Fig. 1, the temperature of the spadix of skunk cabbage is kept at approximately 19°C notwithstanding a fall of the ambient temperature.

The temperature is thus maintained by doubling the respiration rate from
 10 the level of 12°C to that of sub-zero temperature. It has also been considered that the heat production in thermogenic plant species relates to a cellular metabolism called cyanide-non-sensitive/non-phosphorylating electron-transferring pathway, which is controlled by mitochondrial alternative oxidase (AOX) (Berthold and Siedow, 1993; Ito et al., 1997; McIntosh, 1994; Wangner and Krab, 1995).

15

On the other hand, it has been shown that a mitochondrial protein called an uncoupling protein (UCP) plays an important role in generation of heat in mammals. UCP found in the intima of mitochondria make H⁺ flow into the membrane to uncouple aspiration from synthesis of ATP which acts to disperse
 20 chemical energy to metabolic heat (Klaus et al., 1991; Klingenberg and Winkler, 1985; Ricquier et al., 1991). In animals, 3 types of UCPs have been found. UCP1 is primarily distributed in brown adipose tissue (Nichollus and Locke, 1984). UCP2 is found ubiquitously in many tissues (Fleury et al., 1997), and UCP3 is localized specifically in skeletal muscle (Boss et al., 1997).

25

It has been considered that UCPs of mammals, similarly to other carrier proteins of mitochondria, are composed of 6 transmembrane segments, of which the hydrophobic portion is derived from pairing amphipathic α -helix structure (Liu et al., 1988; Maia et al., 1998). It is also known that the activity of these UCPs
 30 decreases depending on purine nucleotides (ATP, GTP, GDP and ADP) attached to

the C-terminal region and increases by free fatty acids (Jezek et al., 1998; Lin and Klingenberg, 1982; Katiyar and Shrago, 1989; Rial et al., 1983; Sluse et al., 1998).

On the contrary, 2 cDNAs encoding UCP-like proteins of plant origin were
5 isolated from potato (StUCP: Laloi et al., 1997) and from *Arabidopsis* (AtPUMP: Maia et al., 1998). Since the expression of StUCP was mainly detected in the flower and the fruit, it has been postulated that StUCP may concern respiration during flowering and maturation of the fruit together with the AOX activity (Laloi et al., 1997).

10

Potato and *Arabidopsis* have been considered to be non-thermogenic plants. However, the expression of StUCP and AtPUMP induced by low temperature. Therefore, it has been suggested that these genes are involved in the heat production (Laloi et al., 1997; Maia et al., 1998).

15

In the thermogenic plants such as skunk cabbage, however, UCP-mediated thermoigenic mechanisms have not yet been identified.

The purpose of the invention of this application is to provide unidentified
20 novel UCP genes derived from a thermogenic plant, skunk cabbage.

The additional purpose of this application is to provide skunk cabbage UCPs which are expression products of the novel genes.

25

Disclosure of Invention

The invention provides thermogenic genes derived from skunk cabbage, i.e., gene SfUCPa of which cDNA comprises the base sequence of SEQ ID NO: 1,
30 and gene SfUCPb of which cDNA comprises the base sequence of SEQ ID NO: 3.

Moreover, the invention provides thermogenic proteins, i.e., protein SfUCPA expressed from SfUCPa, which comprises the amino acid sequence of SEQ ID NO: 2, and protein SfUCPB expressed from SfUCPb, which comprises the amino acid sequence of SEQ ID NO: 4.

In addition, the invention provides cDNA having the base sequence of SEQ ID NO: 1 or a partial sequence thereof, and cDNA having the base sequence of SEQ ID NO: 3 or a partial sequence thereof.

Brief Description of Drawings

Fig. 1 shows the change of the temperature of the spadix in skunk cabbage and that of ambient temperature with a lapse of time.

Fig. 2 shows the results of northern blotting, indicating the expression profile of SfUCPa (A) and SfUCPb (B) in the spadix and leaf of skunk cabbage at room temperature (RT) and during cold treatment (4°C for 3 days). The lower figures respectively show the results of ethidium bromide staining of non-decomposed rRNA.

Fig. 3 compares the alignment of amino acid sequences of SfUCPA and SfUCPB, together with potato UCP (StUCP), *Arabidopsis* UCP (AtPUMP) and human UCP. The asterisks (*) attached under the sequences indicate the same amino acid sequence, and the dot (.) indicates the conservative change in all of the sequences. The boldface indicates the same sequence between SfUCPA and SfUCPB. The gap introduced to optimize the sequence alignment is indicated by a dash (-). The alignment was made using a CLUSTAL W program. The characteristic domains of energy transfer proteins typical of mitochondria are

surrounded by a square. The shaded bars (I~VI) above the upper sequence show estimated transmembrane domains.

Fig. 4 shows a hydrophobic plot of SfUCPA. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM6.

Fig. 5 shows a diagrammatic illustration of SfUCPA topology in the mitochondria membrane.

Fig. 6 shows a hydrophobic plot of SfUCPB. The vertical axis indicates the degree of hydrophobicity and the estimated transmembrane domains are indicated by TM1 to TM4 and TM6.

Fig. 7 shows a diagrammatic illustration of SfUCPB topology in the mitochondria membrane.

Fig. 8 shows the results of *in vitro* translation using respective cDNAs of the genes SfUCPa and SfUCPb as templates. (-) indicates a control, S a sense RNA, and AS an antisense RNA. The asterisk (*) indicates a non-specific product and the empty circle denotes the position of a low molecular translated artificial product synthesized from a small ORF.

Best Mode for Carrying Out the Invention

In the gene SfUCPa of the present invention, its cDNA has the base sequence of SEQ ID NO: 1 and encodes the protein SfUCPA having the amino acid sequence of SEQ ID NO: 2, of which the estimated molecular weight is 32.6 kDa.

In the gene SfUCPb of the present invention, its cDNA (SEQ ID NO: 3) encodes

the protein SfUCPB having the amino acid sequence of SEQ ID NO: 4, of which the estimated molecular weight is 29.0 kDa.

Genes SfUCPa and SfUCPb of the invention are derived from skunk
 5 cabbage, which are expressed specifically in the spadix when the temperature is low. The results of Northern blotting on the total RNAs extracted from skunk cabbage (Ito et al., 1999), confirmed that the expressions of both genes were detected in the spadices but not in the leaves at room temperature (15°C) (Fig. 2). It was also confirmed that the spadix-specific expression of both genes were
 10 induced by cold treatment (4°C for 3 days).

The amino acid sequences of the proteins SfUCPA and SfUCPB that are expressed from the respective genes of the invention have higher homology to the plant UCPs than to the human UCPs (Fig. 3) such that the amino acid sequence of
 15 SfUCPA has homology of 79%, 75%, 44%, 48% and 48% to StUCP, AtPUMP, human UCP, UCP2 and UCP3, respectively. SfUCPB has homology of 71%, 66%, 41%, 43% and 44% to StUCP, AtPUMP, human UCP, UCP2 and UCP3, respectively.

20 In addition, SfUCPA and SfUCPB have high sequence homology (88%) to each other though the region corresponding to the amino acid sequence between the 204th Thr and the 238th Val in SfUCPA is completely deleted in SfUCPB (Fig. 3). Moreover, the 265th Leu of SfUCPA is replaced by Pro in SfUCPB.

25 StUCPA has similar structure to that of other mitochondria UCP proteins. SfUCPA has 6 transmembrane domains as shown by the hydrophobic plot in Fig. 4, of which the topology is as shown in Fig. 5. In addition, this SfUCPA has 3 domains that are characteristic of energy transfer proteins in mitochondria (Fig. 3)(Boss et al., 1997; Maia et al., 1998). On the other hand, SfUCPB is lacking in
 30 the 3rd domain which is characteristic of energy transfer proteins in mitochondrial

(Fig. 3), as well as in the 5th transmembrane domain (Figs. 3 and 6). The topology is located toward the mitochondria matrix at the C-terminal (Fig. 7).

Each protein has a purine nucleotide-binding domain (PNBD) at the C-terminal (Figs. 3, 5 and 7), and it is known that in UCP, binding of the purine nucleotide inhibits the uncoupling function in the mitochondria intima. In SfUCPB, however, there is a possibility that it may have escaped the inhibition of the binding of the purine nucleotide because its C-terminal is located toward the mitochondria matrix. Such a topology has not been found in any UCPs from animals or plants.

The thermogenic genes SfUCPa and SfUCPb provided by the invention are derived from skunk cabbage and are very useful in, for example, development of low temperature-tolerant plants using a genetic recombination technique. The proteins SfUCPA and SfUCPB that are expression products from the above genes are expected as effective components in remedies of diabetes mellitus, obesity, and the like, based on the uncoupling function to ATP synthesis. Moreover, such thermogenic proteins are also promising novel heat generating bio-materials.

The genes SfUCPa and SfUCPb of the invention can be isolated from the genomic DNA of skunk cabbage using the cDNA (SEQ ID NOS: 1 or 3) or a partial sequence thereof of the invention as a probe. For example, a genome library is prepared from the genomic DNA according to a known method. It may be screened by means of colony or plaque hybridization according to a known method using as a probe an oligonucleotide synthesized based on the base sequence of an optional portion of cDNA. Alternatively, the target genetic region may also be identified by means of in situ hybridization for chromosome.

The respective cDNAs of the invention can be cloned, for example, from a cDNA library which is synthesized using a poly(A)⁺RNA of skunk cabbage as a

template. In such a case, an oligonucleotide of an optional portion of cDNA provided by the invention is synthesized, which may be used as a probe to carry out screening by means of colony or plaque hybridization according to a known method. Alternatively, oligonucleotides which can hybridize to both ends of the target cDNA fragment are synthesized, which may be used as primers in preparation of cDNA of the invention by the RT-PCR method from mRNA isolated from the cells of skunk cabbage.

In general, polymorphism is frequently recognized in the genes of eucaryotic cells. In the invention, accordingly, in addition to cDNAs represented by SEQ ID NOS: 1 and 3, those in which one or several nucleotides are added, deleted and/or replaced by (an)other nucleotide(s) in the above cDNA are included. Similarly, proteins in which one or more amino acids are added, deleted and/or replaced by (an)other amino acid(s) due to change of the above nucleotide are also included in the present invention.

In cDNA of the invention, DNA fragments (10bp or more) comprising an optimal part of the base sequences of SEQ ID NOS 1 and 3 are included. In addition, DNA fragments comprising a sense strand or anti-sense strand are also included.

The proteins of the invention, SfUCPA and SfUCPB, may be prepared respectively by a known method, for example, isolation from the spadix of skunk cabbage, preparation by chemical syntheses based on the amino acid sequence provided by the invention, or production by a recombinant DNA technique using cDNA provided by the invention. For example, when the protein is produced by a recombinant DNA technique, RNA is prepared from a vector containing cDNA of the invention by *in vitro* transcription, and this is used as a template for *in vitro* translation to yield the protein. Alternatively, the translational region of cDNA is incorporated into an appropriate expression vector according to a known method,

and the resulting recombinant vector is introduced into *Escherichia coli*, *Bacillus subtilis*, yeast, animal or plant cells. The resulting transformants can be used in expression of the proteins in a large quantity.

5 In the case of the proteins of the invention being produced by *in vitro* translation, the translation region of cDNA of the invention may be incorporated into a vector containing RNA polymerase promotor, and then added to an *in vitro* translation system such as a rabbit reticulocyte lysate or wheat germ extract containing an RNA polymerase corresponding to the promotor. The RNA
10 polymerase promotor is exemplified by T7, T3, SP6, and similar promoters.

 In the case of the proteins of the invention being expressed in a microorganism such as *Escherichia coli*, the translation region of cDNA is incorporated into an expression vector containing an origin replicable in
15 microorganisms, promoter, ribosome binding site, cDNA cloning site, terminator, and the like, to construct a recombinant expression vector, which is then introduced into a host cell and incubated. In this operation, an initiation codon and a stop codon may be added to the front and tail of an optional translation region to obtain a protein fragment containing the optional region. Alternatively,
20 the desired protein may be expressed as a fusion protein with another protein, which may be cleaved with a suitable protease to yield the desired protein. The expression vectors for *Escherichia coli* are exemplified by pUC series, pBluescript II, pET expression system, pGEX expression system, and the like.

25 In the case of the proteins of the invention being expressed in eucaryotic cells, the translation region of cDNA of the invention is incorporated into an expression vector for eucaryotic cells containing a promoter, splicing region, poly(A) additional site, and the like, and introduced into the eucaryotic cells. The expression vector is exemplified by pKA1, pCDM8, pSVK3, pMSG, pSVL,
30 pBK-CMV, pBK-RSV, EVB-vector, pRS, pYES2, and the like. The eucaryotic cells,

many include mammal cultured cells such as monkey renal cell COS7, Chinese hamster ovarian cell CHO, etc., budding yeast, fission yeast, silkworm cell, *Xenopus* egg cell, and the like are commonly used, but not limited thereto. In order to introduce the expression vector into eucaryotic cells, a known method
5 such as electroporation, calcium phosphate method, liposome method, DEAE dextran method, and the like can be utilized.

After expression of the proteins in procaryotic cells or eucaryotic cells according to the aforementioned method, the desired proteins are isolated and
10 purified from the culture in the known combined procedures for separation. For example, treatment with a denaturant (e.g., urea) or surface activator, ultrasonication, digestion with enzymes, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity
15 chromatography, reverse phase chromatography, and the like are involved.

The proteins of the invention, SfUCPA and SfUCPB, also include peptide fragments (5 amino acids or more) involving the optional partial amino acid sequences of SEQ ID NOS: 2 and 4. In addition, the proteins of the invention also
20 include fusion proteins with other optional proteins.

The following examples serve to illustrate the invention of this application specifically in more detail, but are not intended to limit the scope of the invention.

25

Example 1: Cloning of cDNA

The total RNA was extracted from the spadix of skunk cabbage (*Symplocarpus foetidus*) and the complete RNA was determined on 1.0% agarose
30 gel electrophoresis (Ito et al., 1999). Using a mRNA isolation kit (Pharmacia), a

clone associated with the UCP gene family was isolated from the purified poly(A)⁺RNA by RT-PCR. The first strand cDNA was prepared by annealing 20 pmol of cDNA primed primer (5'-TTTTTTTTTTTTTTTTTTTTTTTTTTT-3') into poly(A)⁺RNA (0.1µg), followed by extension with 10 units of reverse transcriptase (New England Biolab) at 37°C for 30 minutes in 20µl of 1×RT buffer containing 10mM 1,4-dithiothreitol and 0.5mM dNTP. The composition of the reaction solution is as follows.

- 10mM Tris-HCl (pH 8.0);
- 50mM KCl;
- 1.5mM MgCl₂;
- 4mM dNTP;
- 0.2 unit of EX Taq polymerase (Takara); and
- 10pmol of two degenerate primers corresponding to the conserved amino acid sequence of the UCP family:

ZF1 (5'-CCIYTIGAYACIGCIAAR-3')

ZR1 (5'-ACWTTCCAISYICCIAWIC-3')

PCR cycle was carried out as follows.

(94°C: 0.5 minute; 50°C: 1 minute; 72°C: 1 minute)×35

Among the PCR products obtained in the above method, the amino acid sequence estimated from the sequence of about 0.8kb cDNA fragment had very high homology to one of the reading frame sequences of the UCP gene family. This fragment, accordingly, was cloned into T-vector (clone p2-1) and used as a probe for library screening.

cDNA (5µg) prepared from the spadix was inserted into λgt11 phage according to the known method (Sambrook et al., 1989) to construct a cDNA library. From this library, 8 clones positive to the above-described probes were isolated and sub-cloned into the pBluescript SK plasmid (Stratagene). From

these clones, clones pz8-1 and pz8-2 were obtained, which respectively contained the full length SfUCPa cDNA and SfUCPb cDNA.

The insert in each clone was sequenced with an auto-sequencer ABI373A using the BcaBest sequencing kit (Takara) and T3, T7 and gene-specific primers. The sequence data were analyzed by means of the GENETYX-Homology Software System version 2.2.0 (Software Development).

cDNA of SfUCPa had the 1,525bp base sequence of SEQ ID NO: 1, and cDNA of SfUCPb had the 2,991bp base sequence of SEQ ID NO: 3. An estimated polyadenylated signal (aataaa) was found upstream of 236bp from the poly(A) sequence in cDNA of SfUCPa, while in cDNA of SfUCPb two polyadenylated sites were recognized at the positions of 1,171bp and 1,243bp. It is noteworthy that cDNA of SfUCPb has a longer 3'-untranslation region than that of SfUCPa.

15

cDNA of SfUCPa had an open reading frame (ORF) encoding 303 amino acids as shown in SEQ ID NO: 1, and this ORF was found to encode the protein SfUCPA of the estimated molecular weight 32.6kDa having the amino acid sequence of SEQ ID NO: 2. On the other hand, cDNA of SfUCPb had an ORF corresponding to 268 amino acids as shown in SEQ ID NO: 3, and found to encode the protein SfUCPB of the estimated molecular weight 29.0kDa.

20

Moreover, it was confirmed from the results of Southern blot analysis that the genome of skunk cabbage contains multiple copies of SfUCPa gene and a single copy of SfUCPb (data not shown).

25

Example 2: *In vitro* translation of cDNA

30

The plasmid clones pz8-1 and pz8-2 obtained in Example 1 were

linearized, on which a sense- or anti-sense RNA was transcribed with T7 RNA polymerase or T3 RNA polymerase according to the protocol of MAXICRIPT transcription kit (Ambion). An equal amount of RNA (4µg) was provided for in vitro translation reaction using a wheat germ extract (Promega) in the presence of ³⁵S-methionine (Amersham). The translation product was analyzed by SDS-PAGE. The gel was fixed, incubated in Amplify (Amersham), then dried, and fluorometrically analyzed.

As a result, it was confirmed that, as shown in Fig. 8, the initiation codon and the stop codon of cDNA isolated in Example 1 functioned successfully since a protein having an expected molecular weight was produced from any of cDNAs only when the sense RNA was used as a template.

Industrial Applicability

As described previously, this application provides novel thermogenic genes SfUCPa and SfUCPb as well as their gene products, i.e., thermogenic proteins SfUCPA and SfUCPB, derived from skunk cabbage (*Symplocarpus foetidus*), and cDNAs used for gene engineering mass production of these proteins. These genes and proteins allow development of low temperature-tolerant plants, development of drugs or methods for treatment of diabetes mellitus or obesity, or development of novel heat generating materials from plants.

References

- Berthold and Siedow (1993) Plant Physiol. 101, 113-119.
- Boss et al. (1997) FEBS Lett. 408, 39-42.
- Fleury et al. (1997) Nature Genetics 15, 269-272.

- Ito, K. et al. (1999) *Plant Sci.* 142, 57-65.
- Ito, K. et al. (1994) *Nucl. Acids Res.* 22, 2036-2041.
- Ito, Y. et al. (1997) *Gene* 12, 121-129.
- Jezek et al. (1998) *Biochem. Biophys. Acta* 1365, 319-327.
- 5 Katiyar and Shrago (1989) *Nati. Acad. Sci. USA* 86, 2559-2562.
- Klaus et al. (1991) *Int. J. Biochem.* 23, 791-810.
- Klingenberg and Winkler (1985) *EMBO J.* 4, 3087-3092.
- Knutson (1974) *Science* 186, 746-747.
- Laloi et al. (1997) *Nature* 389, 135-136.
- 10 Levitt (1980) *Responses of plants to environmental stresses*. 2nd edn. New York: Academic Press.
- Lin and Klingenberg (1982) *Biochemistry* 21, 2950-2956.
- Liu et al. (1988) *J. Cell. Biol.* 107, 503-509.
- Maia et al. (1998) *FEBS Lett.* 429, 403-406.
- 15 McIntosh (1994) *Plant Physiol.* 105, 781-786.
- Nagy et al. (1972) *Science* 178, 1195-1197.
- Nicholls and Locke (1984) *Physiol. Rev.* 64, 1-64.
- Rial et al. (1983) *Eur. J. Biochem.* 137, 197-203.
- Ricquier et al. (1991) *FASEB J.* 5, 2237-2242.
- 20 Sakai and Larcher (1987) *Frost Survival of Plants: Responses and Adaptations to Freezing Stresses*. Berlin and New York; Springer-Verlag.
- Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd eds. New York; Cold Spring Harbor Laboratory Press.
- Schneider and Buchanan (1980) *Amer. J. Bot.* 67, 182-193.
- 25 Sluse et al. (1998) *FEBS Lett.* 433, 237-240.
- Steponkus (1984) *Annu. Rev. Plant Physiol.* 35, 543-581.
- Thomashow (1998) *Plant Physiol.* 118, 1-7.
- Uemura and Steponkus (1997) *Plant Cold Hardiness: Molecular Biology, Biochemistry, and Physiology* (Li and Chen, edn). New York: Plenum Press, pp.
- 30 171-179.

Wagner and Krab (1995) *Physiol. Plant*, 95, 318-325.

CLAIMS

1. A thermogenic gene SfUCPa derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 1.
2. A thermogenic gene SfUCPb derived from skunk cabbage, of which cDNA comprises the base sequence of SEQ ID NO: 3.
3. A thermogenic protein SfUCPA expressed from the gene SfUCPa of Claim 1, which comprises the amino acid sequence of SEQ ID NO: 2.
4. A Thermogenic protein SfUCPB expressed from the gene SfUCPb of Claim 2, which comprises the amino acid sequence of SEQ ID NO: 4.
5. A DNA fragment comprising the base sequence of SEQ ID NO: 1 or a partial sequence thereof.
6. A DNA fragment comprising the base sequence of SEQ ID NO 3: or a partial sequence thereof.

ABSTRACT

The inventions of this application include thermogenic genes named SfUCPa and SfUCPb which are derived from skunk cabbage. cDNA of each gene
5 comprises the base sequence of SEQ ID NO: 1 and 3, respectively. Thermogenic proteins, SfUCPA and SfUCPB, are expressed from genes SfUCPa and SfUCPb, comprises the amino acid sequence of SEQ ID NO: 2 and 4.

SEQUENCE LISTING

<110> Japan Science and Technology Corporation

<120> Plant Thermogenic Genes and Proteins

<150> JP11-167439

<151> 1999-06-14

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 1525

<212> DNA

<213> Symplocarpus foetidus

<220>

<221> CDS

<222> (280).. (1188)

<221> poly A site

<221> (1271).. (1276)

<300>

<301> Ito, K.

<302> Isolation of two distinct cold-inducible cDNAs encoding plant uncoupling proteins from the spadix of skunk cabbage (Symplocarpus foetidus)

<303> Plant Sci.

<304> 149

<305>

<306> 167-173

<307> 1999

<308> GenBank AB024733

<309> 2000-02-25

<400> 1

gaggattcgc agaagaaagg ccagaacccg attccttccc gtcttcttct ccttccgccc 60
aattgcagtt tttcgcagcg gcgtcatcat caagaccctc cgcttttccg cgccaaacgc 120
cttccacccc cacccaatcg ccttccgitt cccgaaatat tcctcttccc tcctcccttt 180
tcttctctac ataaacccta accaccccat cctctcttcc cgcttccgac caccctgcat 240
tctactggga gcccatitga tcgaggtttc ccggcgagg atg ggc gat cac ggc 294

Met Gly Asp His Gly

1

5

ccg agg acc gag atc tcg ttt gcc ggc agt tcg cga gca gca ttc gcc 342
Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser Arg Ala Ala Phe Ala

10

15

20

gct tgc ttc gcc gag ctt tgc acg att ccg ttg gac act gct aaa gtt 390
Ala Cys Phe Ala Glu Leu Cys Thr Ile Pro Leu Asp Thr Ala Lys Val

25

30

35

agg ctt caa ctc caa aag aaa gca gta aca ggt gat gtg gtg gct ttg 438
Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly Asp Val Val Ala Leu

40

45

50

cca aaa tac agg gga atg ttg ggc act gtt gcc act att gcc agg gag 486
Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala Thr Ile Ala Arg Glu

55

60

65

gaa ggt ttg tcg gca ctc tgg aaa gga att gta ccc ggt ttg cat cgt 534
Glu Gly Leu Ser Ala Leu Trp Lys Gly Ile Val Pro Gly Leu His Arg

70

75

80

85

caa tgc ctc ttt gga ggg cta cga att ggg ttg tat gaa cca gtt aag 582
Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu Tyr Glu Pro Val Lys

90

95

100

tcc ttt tat gtt gga gat aac ttt gtt gga gat att cct tta tcc aag 630
Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp Ile Pro Leu Ser Lys

105

110

115

aaa ata ctt gct ggg ctt aca aca ggt gca tta gca att ata gtg gca 678
 Lys Ile Leu Ala Gly Leu Thr Thr Gly Ala Leu Ala Ile Ile Val Ala
 120 125 130
 aat ccc act gac ctt gtt aaa gtt cga ctt caa tct gaa ggt aaa ctc 726
 Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln Ser Glu Gly Lys Leu
 135 140 145
 ccc cct ggg gta ccg aga cgt tat tca ggg gcg cta aat gct tat tca 774
 Pro Pro Gly Val Pro Arg Arg Tyr Ser Gly Ala Leu Asn Ala Tyr Ser
 150 155 160 165
 acc ata gtc aaa aag gaa gga ctt ggt gct ctg tgg act ggg ctt ggt 822
 Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu Trp Thr Gly Leu Gly
 170 175 180
 cct aat att gcc cgc aat gct att ata aat gct gct gaa ttg gcc agt 870
 Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala Ala Glu Leu Ala Ser
 185 190 195
 tat gat caa gtg aaa cag aca atc tta aaa tta cca gga ttc tca gat 918
 Tyr Asp Gln Val Lys Gln Thr Ile Leu Lys Leu Pro Gly Phe Ser Asp
 200 205 210
 aat att ttt act cat att tta gcc ggt ctg ggg gca ggt ttt ttt gcc 966
 Asn Ile Phe Thr His Ile Leu Ala Gly Leu Gly Ala Gly Phe Phe Ala
 215 220 225
 gtc tgt atc ggt tct cct gtt gat gtg atg aag tct aga atg atg gga 1014
 Val Cys Ile Gly Ser Pro Val Asp Val Met Lys Ser Arg Met Met Gly
 230 235 240 245
 gat tca gcc tac aaa agc aca ttt gat tgt ttc atc aag aca ttg aaa 1062
 Asp Ser Ala Tyr Lys Ser Thr Phe Asp Cys Phe Ile Lys Thr Leu Lys
 250 255 260
 aat gat ggg ctt ctt gct ttt tac aag ggg ttt atc cca aac ttt ggt 1110

Asn Asp Gly Leu Leu Ala Phe Tyr Lys Gly Phe Ile Pro Asn Phe Gly
 265 270 275
 cgg tta gga tcg tgg aat gtg atc atg ttt ttg acc ttg gag cag gtc 1158
 Arg Leu Gly Ser Trp Asn Val Ile Met Phe Leu Thr Leu Glu Gln Val
 280 285 290
 aag aag ttt ttc atc aaa gag gtg cca aat taatacattg aactcggata 1208
 Lys Lys Phe Phe Ile Lys Glu Val Pro Asn
 295 300
 ggagtagaaa gaaagggttt ttgtggaatt ttctctaccg gtgtggatcc tggcgagaga 1268
 caaataaadc ttctgactg ctgagatgtg tacctttttt atgaatggtt cttttcttat 1328
 agaggacaga gaaaagaaaa aaaaaatcat tgtcatttac tctttttccc catttctgct 1388
 gctaactctg gtaggagaag aaaagtctta cattgagtga taacgttggt ctctgcatcc 1448
 attatttttc agagatacta ttgacacat gaaaagtaat gcacatcagg ttttttttaa 1508
 aaaaaaaaaa aaaaaaa 1525

<210> 2

<211> 303

<212> PRT

<213> Symplocarpus foetidus

<400> 2

Met Gly Asp His Gly Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser
 1 5 10 15
 Arg Ala Ala Phe Ala Ala Cys Phe Ala Glu Leu Cys Thr Ile Pro Leu
 20 25 30
 Asp Thr Ala Lys Val Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly
 35 40 45
 Asp Val Val Ala Leu Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala
 50 55 60

Thr Ile Ala Arg Glu Glu Gly Leu Ser Ala Leu Trp Lys Gly Ile Val
 65 70 75 80
 Pro Gly Leu His Arg Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu
 85 90 95
 Tyr Glu Pro Val Lys Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp
 100 105 110
 Ile Pro Leu Ser Lys Lys Ile Leu Ala Gly Leu Thr Thr Gly Ala Leu
 115 120 125
 Ala Ile Ile Val Ala Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln
 130 135 140
 Ser Glu Gly Lys Leu Pro Pro Gly Val Pro Arg Arg Tyr Ser Gly Ala
 145 150 155 160
 Leu Asn Ala Tyr Ser Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu
 165 170 175
 Trp Thr Gly Leu Gly Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala
 180 185 190
 Ala Glu Leu Ala Ser Tyr Asp Gln Val Lys Gln Thr Ile Leu Lys Leu
 195 200 205
 Pro Gly Phe Ser Asp Asn Ile Phe Thr His Ile Leu Ala Gly Leu Gly
 210 215 220
 Ala Gly Phe Phe Ala Val Cys Ile Gly Ser Pro Val Asp Val Met Lys
 225 230 235 240
 Ser Arg Met Met Gly Asp Ser Ala Tyr Lys Ser Thr Phe Asp Cys Phe
 245 250 255
 Ile Lys Thr Leu Lys Asn Asp Gly Leu Leu Ala Phe Tyr Lys Gly Phe
 260 265 270
 Ile Pro Asn Phe Gly Arg Leu Gly Ser Trp Asn Val Ile Met Phe Leu
 275 280 285

Thr Leu Glu Gln Val Lys Lys Phe Phe Ile Lys Glu Val Pro Asn

290

295

300

<210> 3

<211> 2991

<212> DNA

<213> *Symplocarpus foetidus*

<220>

<221> CDS

<222> (286).. (1089)

<221> poly A site

<222> (1171) (1176)

<221> poly A site

<222> (1243) (1248)

<300>

<301> Ito, K.

<302> Isolation of two distinct cold-inducible cDNAs encoding plant uncoupling proteins from the spadix of skunk cabbage (*Symplocarpus foetidus*)

<303> Plant Sci.

<304> 149

<305>

<306> 167-173

<307> 1999

<308> GenBank AB024734

<309> 2000-02-25

<400> 3

tggtggtgac gaggtagcag gattcgagcaga agaaaggcca gaacccgatt ccttcccgtc 60

ttcttctcct tccgcccaat tgcagttttt cgcagcgggt catcatcaag accctccgcc 120

tttccgcgcc aaacgccttc cacccaatcc ctccgtttcc cgaaatatcc cccttccttc 180
 ccttttcttc tctacataaa ccctaaccac ccccatcctc tcctcccgct tccgaccacc 240
 ctgcattcta ctgggatccc atttgatcga cgtttcccg gagg atg ggc gat cac 297

Met Gly Asp His

1

ggc ccg agg acc gag atc tcg ttt gcc ggc agt tcg cga gca gca ttc 345
 Gly Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser Arg Ala Ala Phe

5 10 15 20

gcc gct tgc ttc gcc gag ctc tgt acg att ccg ttg gac act gct aaa 393
 Ala Ala Cys Phe Ala Glu Leu Cys Thr Ile Pro Leu Asp Thr Ala Lys

25 30 35

gtt agg ctt cag ctc caa aag aaa gca gta aca ggt gat gtg gtg gct 441
 Val Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly Asp Val Val Ala

40 45 50

ttg cca aaa tac agg gga atg ttg ggc act gtt gcc act att gcc agg 489
 Leu Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala Thr Ile Ala Arg

55 60 65

gag gaa ggt ttg tcg gca ctc tgg aaa gga att gta ccc ggt ttg cat 537
 Glu Glu Gly Leu Ser Ala Leu Trp Lys Gly Ile Val Pro Gly Leu His

70 75 80

cgt caa tgc ctc ttt gga ggg cta cga att ggg ttg tat gaa cca gtt 585
 Arg Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu Tyr Glu Pro Val

85 90 95 100

aag tcc ttt tat gtt gga gat aac ttt gtt gga gat att cct tta tcc 633
 Lys Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp Ile Pro Leu Ser

105 110 115

aag aaa ata ctt gct ggg ctt aca aca ggt gca tta gca att ata gtg 681
 Lys Lys Ile Leu Ala Gly Leu Thr Thr Gly Ala Leu Ala Ile Ile Val

120	125	130	
gca aat ccg act gac ctt gtt aaa gtt cga ctt caa tct gaa ggt aaa			729
Ala Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln Ser Glu Gly Lys			
135	140	145	
ctc ccc cct ggg gta cca aga cgt tat tca ggg gcg cta aat gct tat			777
Leu Pro Pro Gly Val Pro Arg Arg Tyr Ser Gly Ala Leu Asn Ala Tyr			
150	155	160	
tca acc ata gtc aaa aag gaa gga ctt ggt gct ctg tgg act ggg ctt			825
Ser Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu Trp Thr Gly Leu			
165	170	175	180
ggg cct aat att gcc cgc aat gct att ata aat gct gct gaa ttg gcc			873
Gly Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala Ala Glu Leu Ala			
185	190	195	
agt tat gat caa gtg aaa cag atg aag tct aga atg atg gga gat tca			921
Ser Tyr Asp Gln Val Lys Gln Met Lys Ser Arg Met Met Gly Asp Ser			
200	205	210	
gcc tac aaa agc aca ttt gat tgt ttc atc aag acg ttg aaa aat gat			969
Ala Tyr Lys Ser Thr Phe Asp Cys Phe Ile Lys Thr Leu Lys Asn Asp			
215	220	225	
ggg cct ctt gct ttt tac aag ggg ttt atc cca aac ttt ggt cgg tta			1017
Gly Pro Leu Ala Phe Tyr Lys Gly Phe Ile Pro Asn Phe Gly Arg Leu			
230	235	240	
gga tcg tgg aat gtg atc atg ttt ttg acc ttg gag cag gtc aag aag			1065
Gly Ser Trp Asn Val Ile Met Phe Leu Thr Leu Glu Gln Val Lys Lys			
245	250	255	260
ttc ttc atc aaa gag gtg cca aat taatacattg aagtcggata ggagtagaaa			1119
Phe Phe Ile Lys Glu Val Pro Asn			

aaaagggttt ttgtggaatt ttctctaccg gtgtggatcc tggcgagaga gaataaatct 1179
tcctgactgc tcagatgttg tacctttttt atgaatggtt cttttcttat agaggacaga 1239
gaaaaataaaa gaaaaattca ttgtcatgta ctctttttcc ccatttctgc tgagtagcag 1299
ctataccaag cagactttgt tgcttggctg ctgctaactt thtagctgaa gaaaagtctt 1359
acattgagtg ataacgttgt tctctgcatc cattattttt cagagttact atttgacaca 1419
tgaaaagttt tttttttttt tttttttttt aacaggcagc aaatagagga atcgatctca 1479
cgactatcct ctttattcat taacaggcat acaaacttag ggagagcatg cagggtatat 1539
atcaaaatat acccttttat tagacatttt gcgtacacag ttggtcctca aacgactgta 1599
tctagcagcc aatttttaga ccacattaag acagagagaa acaagcagaa gaacagggta 1659
ccatacatat ataggtaata attaagatga tgaacatagc ataggttcat gatctacttc 1719
ttcttcacgt acacatgatg caccagctga atgggaatct tggtcacat atggcatgaa 1779
agtacgtcat gtgcagacgt tatatagtgt tcttcttacc attcagcagc agcaccagag 1839
gcatcaaaca ctggggctct gacagggtat gagggttaca ttgcgatccc acacatgccg 1899
tagggagtca cattgcgttt gatttttatg tatccagcct gtccccacct agtgccccat 1959
gagttcctta caagccaata atccttccca tcctccgaac catacctat tatcaccaca 2019
gcatggtcga tacgttgacc acatgggtcca gcaaatacgc ccgagggtga gtgttggaat 2079
ccagcgcccg aagcctcaag agcaacactg acagggtgct ttgcgactgc atactgtagg 2139
ctaacctgt tgtacggaga aacattttca tacgcatcaa tcgaggtgac ttttaataaga 2199
tttgctctgc aagttcctcg acgtcctgtg tacgggtaat ttttggaaaa aggagaaggg 2259
cgggagaag cgcgctctc tgctcgcgac gggttaattc ttcatatggc cacttcgagc 2319
atggcttcgg cagcctccag ctctgttctc actccggcct cccctccacc accacgccgg 2379
ttccccgctc cgctttcttt ccagggtcgc caggaggctc gtggtggtgc gggccgagga 2439
agccgcgacg acccccgcgc ccggcgccgg cggaggagc cgcgccgccg cccccaagcc 2499
gccaccgatc gggcccaaga gggggtcaaa gttgatata tagttcttaa tttctttccc 2559
gtgtggcttc tggagttaga ttgttttctt ctctctttt tttgtttgt tttttcaatt 2619
taaattttat tctcatctgt ggacgacctt ccatcgggtt tttcgtccct ctgcaggtg 2679
aagatctccg gaaggaatcc tactggttca acggtgtcgg atcggtggtg gctgttgatc 2739
aggatccggc gactcgatac ccggtcgtgg ttcggttcac caaggtcaac tatgcgaacg 2799

tctcgaccaa caactacgca ctggacgaga tcctggaggt gaaatgaggg tcggcgggcg 2859
 tggtcggtcg ggcatgtcac gatgatgtat tttcgcagtt ggtagtgtaa aataccatgt 2919
 cattcgtgta aaactctttc gttcgccaaa tcctcagttg aaattttaat tcccagccag 2979
 taaaaaaaa aa 2991

<210> 4

<211> 268

<212> PRT

<213> Symplocarpus foetidus

<400> 4

Met Gly Asp His Gly Pro Arg Thr Glu Ile Ser Phe Ala Gly Ser Ser

1 5 10 15

Arg Ala Ala Phe Ala Ala Cys Phe Ala Glu Leu Cys Thr Ile Pro Leu

20 25 30

Asp Thr Ala Lys Val Arg Leu Gln Leu Gln Lys Lys Ala Val Thr Gly

35 40 45

Asp Val Val Ala Leu Pro Lys Tyr Arg Gly Met Leu Gly Thr Val Ala

50 55 60

Thr Ile Ala Arg Glu Glu Gly Leu Ser Ala Leu Trp Lys Gly Ile Val

65 70 75 80

Pro Gly Leu His Arg Gln Cys Leu Phe Gly Gly Leu Arg Ile Gly Leu

85 90 95

Tyr Glu Pro Val Lys Ser Phe Tyr Val Gly Asp Asn Phe Val Gly Asp

100 105 110

Ile Pro Leu Ser Lys Lys Ile Leu Ala Gly Leu Thr Thr Gly Ala Leu

115 120 125

Ala Ile Ile Val Ala Asn Pro Thr Asp Leu Val Lys Val Arg Leu Gln

130 135 140

Ser Glu Gly Lys Leu Pro Pro Gly Val Pro Arg Arg Tyr Ser Gly Ala
145 150 155 160
Leu Asn Ala Tyr Ser Thr Ile Val Lys Lys Glu Gly Leu Gly Ala Leu
165 170 175
Trp Thr Gly Leu Gly Pro Asn Ile Ala Arg Asn Ala Ile Ile Asn Ala
180 185 190
Ala Glu Leu Ala Ser Tyr Asp Gln Val Lys Gln Met Lys Ser Arg Met
195 200 205
Met Gly Asp Ser Ala Tyr Lys Ser Thr Phe Asp Cys Phe Ile Lys Thr
210 215 220
Leu Lys Asn Asp Gly Pro Leu Ala Phe Tyr Lys Gly Phe Ile Pro Asn
225 230 235 240
Phe Gly Arg Leu Gly Ser Trp Asn Val Ile Met Phe Leu Thr Leu Glu
245 250 255
Gln Val Lys Lys Phe Phe Ile Lys Glu Val Pro Asn
260 265

Fig. 1




Fig.2




Fig.3 Top

I

SfUCPA	1	--MGDHGPRTEISFAGSSR-AAFAACFAELCT	PLDTAKVRLQLQKKA	VDV-VALPKY	56
SfUCPB	1	--MGDHGPRTEISFAGSSR-AAFAACFAELCT	PLDTAKVRLQLQKKA	VDV-VALPKY	56
StUCP	1	MGGGDHGGKSDISFAGIFASSAFACFAEACT	PLDTAKVRLQLQKKA	VEGCG-LALPKY	59
AtPUMP	1	---MVAAGKSOLSLPKTFACSAFAACVGEVCT	PLDTAKVRLQLQKSA	FTLAGDVTLPKY	57
human UCP1	1	-MGGLTASDVHPTLGVQLFSAPIAACLADVI	THPLDTAKVRLQVQ	GECP----TSSVIRY	55
human UCP2	1	-MVGFKATDVPPPTATVKFLGAGTAACIADLI	THPLDTAKVRLQIQGESQ	GPVPRATASAQY	59
human UCP3	1	-MVGLKPSOVPPPTMAVKFLGAGTAACFADLV	THPLDTAKVRLQIQGENQ	-AVQTARLVQY	58

II

SfUCPA	57	RGMLGTVATIAREEGLSALWKGI	VPGLHRQCLFGGLRIGLYEPVKS	FYVG--DNFVGDIP	114
SfUCPB	57	RGMLGTVATIAREEGLSALWKGI	VPGLHRQCLFGGLRIGLYEPVKS	FYVG--DNFVGDIP	114
StUCP	60	RGLLGTVGTIAKEEGLASLWKGI	VPGLHRQCIYGGLRIGMYEPVKNL	YVG--KDHVGDVP	117
AtPUMP	58	RGLLGTVGTIAREEGLSRLWKGV	VPGLHRQCLFGGLRIGMYEPVKNL	YVFTGKDFVGDVP	117
human UCP1	56	KGVLTGITAVVKTEGRMKLYSGL	PAGLQRQISSASLRIGLYD	TVQEFFLTA---GKETAPS	112
human UCP2	60	RGVMGTILTMVRTEGPRSLYNG	LVAGLQRQMSFASVRIGLYDS	VKKQFYT----KGSEHAS	115
human UCP3	59	RGVLGTILTMVRTEGPCSPYNG	LVAGLQRQMSFASIRIGLYDS	VKKQVYTF----KGADNSS	115

Fig.3 Middle

III

SfUCPA 115 LSKKILAGLTGALAIIVANPTDLVKVRLQSEGKLPAG--VPRRYSGALNAY--STIVKKE 171
 SfUCPB 115 LSKKILAGLTGALAIIVANPTDLVKVRLQSEGKLPAG--VPRRYSGALNAY--STIVKKE 171
 StUCP 118 LSKKILAGLTGALGITIANPTDLVKVRLQAECKLPAG--VPRRYSGALNAY--STIVKQE 174
 AtPUMP 118 LSKKILAGLTGALGIMVANPTDLVKVRLQAECKLAAG--APRRYSGALNAYFTSTIVRQE 176
 human UCP1 113 LGSKILAGLTGGVAVFIGQPTLVVKVRLQAQSHLHG--IKPRYTGTYNAY--RIIATTE 168
 human UCP2 116 IGSRLLAGSTTGALAVAVACPTDVVKVRFQAARAG---GRRYQSTVNAY--KTIAREE 170
 human UCP3 116 LTTRILAGCTTGAMAVTCAQPTDVVKVRFQASIHLPSSDRKYSGTMDAY--RTIAREE 173
 ** * * *

IV

V

SfUCPA 172 GLGALWTGLGPNLARNAIINAAELASYDQVKQTILKLPFSDNIFTHILAG--LGAGFFA 229
 SfUCPB 172 GLGALWTGLGPNLARNAIINAAELASYDQVKQ----- 203
 StUCP 175 GVRALWTGLGPNIGRNAIINAAELASYDQVKEAVLRIPGFTDNVTVTHLIAG--LGAGFFA 232
 AtPUMP 177 GVRALWTGLGPNVARNAIINAAELASYDQVKETILKIPGFTDNVTVTHILSGLFTGAGFFA 236
 human UCP1 169 GLTGLWKGTTPNLRSVIINCTELVTYDLMKEAFVKNNILADVPCHLVSA--LIAGFCA 226
 human UCP2 171 GFRGLWKGTSPNVARNAIVNCAELVTYDLIKDALLLKANLMTDOLPCHFTSA--FGAGFCT 228
 human UCP3 174 GVRGLWKGTLPNMRNAIVNCAEVVTTYDILKEKLLOYHLLTDNFPCHFVSA--FGAGFCA 231
 * ** * * *

Fig. 3 Down

VI

230	VCIGSPVDVVKSRMAGDS--AYKSTFDCF	IKILKNDGLLAFYKGFIPNFGRLG--SWNV	285
204	-----MKSRMAGDS--AYKSTFDCF	IKILKNDGPLAFYKGFIPNFGRLG--SWNV	250
233	VCIGSPVDVVKSRMAGDS--AYKNTLDCFVK	TLKNDGPLAFYKGFIPNFGRLG--SWNV	288
237	VCIGSPVDVVKSRMAGDSG--AYKGTIDCFVK	TLKSDGPMFAYKGFIPNFGRLGSTWNV	295
227	TAMSSPVDVVKTRHINSPPGQYKSVFNCAMKV	FTNEGPTAFKGLVPFSLRLG--SWNV	284
229	TVIASPVDVVKTRHMNSALGQYSSAGHCAL	TMLQKEGPRAFYKGFMPFSLRLG--SWNV	286
232	TVVASPVDVVKTRHMNSPPGQYFSPLDCMIKM	VAQEGPTAFYKGFTPFSLRLG--SWNV	289

	<u>PNBD</u>	
SfUCPA	286 MFLLTLEQVKKFFIKEVPN-----	303
SfUCPB	251 MFLLTLEQVKKFFIKEVPN-----	268
StUCP	289 MFLLTLEQAKKFVKSLSP-----	306
AtPUMP	296 MFLLTLEQAKKYVRELDASKRN--	316
human UCP1	285 MFVCFEQLKRELSKSRQTHDCAT	307
human UCP2	289 MFVTYTEQLKRALMAACTSREAPF	309
human UCP3	290 MFVTYTEQLKRALKVKVQMLRESPP	312

Fig.4




Fig.5




Fig.6




Fig. 7





Fig.8



Rev. 3-21-01
DECLARATION AND POWER OF ATTORNEY FOR U.S. PATENT APPLICATION

(X) Original () Supplemental () Substitute (X) PCT () DESIGN

As below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that I verily believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Title: PLANT THERMOGENIC GENES AND PROTEINS

of which is described and claimed in:

() the attached specification, or

(X) the specification in application Serial No. _____, filed December 14, 2001, and with amendments through _____, or

(X) the specification in International Application No. PCT/JP00/03806, filed June 12, 2000, and as amended on _____ (if applicable).

I hereby state that I have reviewed and understand the content of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge my duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim priority benefits under Title 35, United States Code, §119 (and §172 if this application is for a Design) of any application(s) for patent or inventor's certificate listed below and have also identified below any application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

COUNTRY	APPLICATION NO.	DATE OF FILING	PRIORITY CLAIMED
Japan	1999-167439	June 14, 1999	Yes

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

APPLICATION SERIAL NO.	U.S. FILING DATE	STATUS: PATENTED, PENDING, ABANDONED

And I hereby appoint Michael R. Davis, Reg. No. 25,134; Matthew M. Jacob, Reg. No. 25,154; Warren M. Cheek, Jr., Reg. No. 33,367; Nils Pedersen, Reg. No. 33,145; Charles R. Waus, Reg. No. 33,142; and Michael S. Huppert, Reg. No. 40,268, who together constitute the firm of WENDEROTH, LIND & PONACK, L.L.P., as well as any other attorneys and agents associated with Customer No. 000513, to prosecute this application and to transact all business in the U.S. Patent and Trademark Office connected therewith.

I hereby authorize the U.S. attorneys and agents named herein to accept and follow instructions from NISHIZAWA & ASSOCIATES as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and myself. In the event of a change in the persons from whom instructions may be taken, the U.S. attorneys named herein will be so notified by me.

Direct Correspondence to Customer No:



000513

PATENT TRADEMARK OFFICE

Direct Telephone Calls to:

WENDEROTH, LIND & PONACK, L.L.P.
2033 "K" Street, N.W., Suite 800
Washington, D.C. 20006-1021

Phone: (202) 721-8200
Fax: (202) 721-8250

Full Name of
First Inventor

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

ITO

Kikukatsu

Residence &
Citizenship

CITY

STATE OR COUNTRY

COUNTRY OF CITIZENSHIP

Iwate

Japan

JPX

Japan

Post Office
Address

ADDRESS

CITY

STATE OR COUNTRY

ZIP CODE

Estate Midoriga-oka 207, 17-7, Midoriga-oka 1-chome, Morioka-shi, Iwate,
Japan

Full Name of
Second Inventor

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

Residence &
Citizenship

CITY

STATE OR COUNTRY

COUNTRY OF CITIZENSHIP

Post Office
Address

ADDRESS

CITY

STATE OR COUNTRY

ZIP CODE

Full Name of
Third Inventor

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

Residence &
Citizenship

CITY

STATE OR COUNTRY

COUNTRY OF CITIZENSHIP

Post Office
Address

ADDRESS

CITY

STATE OR COUNTRY

ZIP CODE

Full Name of
Fourth Inventor

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

Residence &
Citizenship

CITY

STATE OR COUNTRY

COUNTRY OF CITIZENSHIP

Post Office
Address

ADDRESS

CITY

STATE OR COUNTRY

ZIP CODE

Full Name of
Fifth Inventor

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

Residence &
Citizenship

CITY

STATE OR COUNTRY

COUNTRY OF CITIZENSHIP

Post Office
Address

ADDRESS

CITY

STATE OR COUNTRY

ZIP CODE

Full Name of
Sixth Inventor

FAMILY NAME

FIRST GIVEN NAME

SECOND GIVEN NAME

Residence &
Citizenship

CITY

STATE OR COUNTRY

COUNTRY OF CITIZENSHIP

Post Office Address	ADDRESS	CITY	STATE OR COUNTRY	ZIP CODE
------------------------	---------	------	------------------	----------

I further declare that all statements made herein of my own knowledge are true, and that all statements on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1st Inventor Kikukatsu ITO Date January 11, 2002
 Kikukatsu ITO
 2nd Inventor _____ Date _____
 3rd Inventor _____ Date _____
 4th Inventor _____ Date _____
 5th Inventor _____ Date _____
 6th Inventor _____ Date _____

The above application may be more particularly identified as follows:

U.S. Application Serial No. _____ Filing Date December 14, 2001

Applicant Reference Number 99-F-071US/YS Atty Docket No. 2001 1838A

Title of Invention PLANT THERMOGENIC GENES AND PROTEINS